



Unretouched photos taken at maximum frown before treatment with BOTOX® Cosmetic and taken at maximum frown after treatment with BOTOX® Cosmetic at day 7. The photos featured here are of an actual BOTOX® Cosmetic patient. Individual results may vary.

In clinical trials at day 7, 74% of patients demonstrated *none* or *mild* glabellar line severity at maximum frown as compared to 6% in placebo; at day 30, 80% of patients demonstrated the same as compared to 3% in placebo.¹

Side effects associated with the injection include localized pain, infection, inflammation, tenderness, swelling, redness, and/or bleeding/bruising.

1. BOTOX® Cosmetic Prescribing Information, August 2009.

Indication

BOTOX® Cosmetic is a prescription medicine that is injected into muscles and used to improve the look of moderate to severe frown lines between the eyebrows (glabellar lines) in people 18 to 65 years of age for a short period of time (temporary).

IMPORTANT SAFETY INFORMATION

BOTOX® Cosmetic may cause serious side effects that can be life threatening. Call your doctor or get medical help right away if you have any of these problems any time (hours to weeks) after injection of BOTOX® Cosmetic:

- **Problems swallowing, speaking, or breathing**, due to weakening of associated muscles, can be severe and result in loss of life. You are at the highest risk if these problems are pre-existing before injection. Swallowing problems may last for several months
- **Spread of toxin effects.** The effect of botulinum toxin may affect areas away from the injection site and cause serious symptoms including: loss of strength and all-over muscle weakness, double vision, blurred vision and drooping eyelids, hoarseness or change or loss of voice (dysphonia), trouble saying words clearly (dysarthria), loss of bladder control, trouble breathing, trouble swallowing

There has not been a confirmed serious case of spread of toxin effect when BOTOX® Cosmetic has been used at the recommended dose to treat frown lines.

The dose of BOTOX® Cosmetic is not the same as, or comparable to, another botulinum toxin product.

Serious and or immediate allergic reactions have been reported. These reactions include itchy rash, swelling, and shortness of breath. Tell your doctor or get medical help right away if you experience any such symptoms; further injection of BOTOX® Cosmetic should be discontinued.

Do not take BOTOX® Cosmetic if you: are allergic to any of the ingredients in BOTOX® Cosmetic (see Medication Guide for ingredients); had an allergic reaction to any other botulinum toxin

product such as *Myobloc*® or *Dysport*®; have a skin infection at the planned injection site.

Tell your doctor about all your muscle or nerve conditions, such as amyotrophic lateral sclerosis [ALS or Lou Gehrig's disease], myasthenia gravis, or Lambert-Eaton syndrome, as you may be at increased risk of serious side effects including severe dysphagia (difficulty swallowing) and respiratory compromise (difficulty breathing) from typical doses of BOTOX® Cosmetic.

Tell your doctor about all your medical conditions, including if you have: plans to have surgery; had surgery on your face; weakness of forehead muscles, such as trouble raising your eyebrows; drooping eyelids; any other abnormal facial change; are pregnant or plan to become pregnant (it is not known if BOTOX® Cosmetic can harm your unborn baby); are breast-feeding or plan to breast-feed (it is not known if BOTOX® Cosmetic passes into breast milk).

Tell your doctor about all the medicines you take, including prescription and nonprescription medicines, vitamins, and herbal products.

BOTOX® Cosmetic may cause loss of strength or general muscle weakness, or vision problems. **If this happens, do not drive a car, operate machinery, or do other dangerous activities.**

Other side effects of BOTOX® Cosmetic include: dry mouth, discomfort or pain at the injection site, tiredness, headache, neck pain, and eye problems: double vision, blurred vision, decreased eyesight, drooping eyelids, swelling of your eyelids, and dry eyes.

For more information refer to the Medication Guide or talk with your doctor.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Full Product Information, including Medication Guide, has been provided to your doctor.

Dysport is a registered trademark of Ipsen Biopharm Limited. *Myobloc* is a registered trademark of Solstice Neurosciences, Inc.